

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

REMARKS

The examiner has issued a restriction requirement in the present case. Applicants provisionally elect species 1, as defined by the examiner, and have introduced claims 14 to 25 as drawn to this species. Additionally, applicants traverse the restriction requirement, and request that the examiner reconsider in light of the following.

Applicants bring to the examiner's attention the nature of the present invention, and submit that as it has passed the international phase of PCT prosecution, the invention has been determined to have unity of invention by the examiner in that phase. It is hoped that this determination will influence the present examiner to withdraw the restriction requirement that has been made.

The examiner has stated that species Ia1.1-1.456 ("1"), Ib1.1-1.456 ("2") and 2.1-3.6 ("3") do not form a single general inventive concept under PCT Rule 13.1. The species are said to lack the same or corresponding special technical features for the reason that "[they] lack a common core" (office action, pp. 2-3). As the examiner makes no further discussion of this allegedly missing feature, and as the species groupings seem to contradict the intuitive understanding that is most directly derived from this assertion, applicants respectfully submit that the restriction requirement does not properly follow PCT Rule 13.1 or 37 CFR §1.475(a). The missing "common core" is not clearly indicated, and each of the groupings shares a special technical feature that defines it generically over the prior art.

To indicate the applicants' clouded understanding of the "common core" set forward by the examiner, applicants first note that the commonality extant within species 3 (i.e., that commonality between 2.1-2.66 and 3.1-3.6) is formula I of the presently claimed invention. The difference between species 1 and 2, is in the group  $R^4$  of formula I, and these two species also, then, share the commonality of this formula. The special technical feature that defines these species collectively over the prior art is this basic formula I, in which  $R^4$  is one of two main formulae (IIa and IIb) that thereby create two groupings, species 1 (Ia1.1-1.456) and a portion of species 3 (2.1-2.66), and species 2 (Ib1.1-1.456) and another portion of species 3 (3.1-3.6).

Formula I, with its two iterations, has been used by the examiner as both a basis for separation of species 1 and 2, and as a basis for inclusion of an equally disparate set of compounds in species 3. The examiner's statement that no "common core" exists between all three species seems convenient, at the least, and not demonstrably in compliance with PCT Rule 13 or 37 CFR §1.475(a). As applicants assert that formula I is a special technical feature defining the three groupings collectively over the prior art, they additionally request that unity of invention be recognized in the present case, and that the restriction requirement be withdrawn.

In view of the foregoing amendments and remarks, applicants consider that the rejections of record have been obviated and respectfully solicit passage of the application to issue.


**Please find attached a check for \$90.00 for the additional claims fee.**

WITSCHER et al.

Serial No. 09/763,704

Please charge any shortage in fees due in connection with the filing of this paper, including Extension of Time fees to Deposit Account No. 11-0345. Please credit any excess fees to such deposit account.

Respectfully submitted,  
KEIL & WEINKAUF

A handwritten signature in black ink, appearing to read "David C. Liechty", written over a horizontal line.


David C. Liechty  
Reg. No. P-48,692

1101 Connecticut Ave., N.W.  
Washington, D.C. 20036  
(202)659-0100  
DCL/mks

**VERSION WITH MARKINGS TO SHOW CHANGES MADE TO THE CLAIMS**


Please amend claim 6-9 as follows:

- 6.(amended) A process for preparing compounds of the formula I as claimed in claim 1 where  $R^5$  = halogen, which comprises reacting a cyclohexanedione derivative of the formula III,




where the variables  $R^1$  to  $R^3$ ,  $R^6$  and  $[1]$  are each as defined in claim 1, with a halogenating agent.

- 7.(amended) A process for preparing compounds of the formula I as claimed in claim 1 where  $R^5$  =  $OR^7$ ,  $OSO_2R^8$ ,  $OPR^8R^9$ ,  $OPOR^8R^9$  or  $OPSR^8R^9$ , which comprises reacting a cyclohexanedione derivative of the formula III,




where the variables  $R^1$  to  $R^3$ ,  $R^6$  and  $[1]$  are each as defined in claim 1, with a compound of the formula  $IV\alpha$ ,  $IV\beta$ ,  $IV\gamma$ ,  $IV\delta$  or  $IV\epsilon$ ,



where the variables  $R^7$  to  $R^9$  are each as defined in claim 1 and  $L^1$  is a nucleophilically replaceable leaving group.


8.(amended) A process for preparing compounds of the formula I as claimed in claim 1

where  $R^5 = OR^7, SR^7, POR^8R^9, NR^{10}R^{11}, ONR^{11}R^{12}$ , N-linked heterocyclyl or O-(N-linked heterocyclyl), which comprises reacting a compound of the formula I  $\alpha$  ( $\equiv$  I where  $R^5 = \text{halogen, } OSO_2R^8$ ),



I where  $R^5 = \text{halogen or } OSO_2R^8$


where the variables  $R^1$  to  $R^3$ ,  $R^6$  and  $[1]$  are each as defined in claim 1, with a compound of the formula  $V\alpha, V\beta, V\gamma, V\delta, V\epsilon, V\eta, V\theta$ ,



where the variables  $R^7$  to  $R^{12}$  are each as defined in claim 1, if appropriate in the presence of a base.

9.(amended) A process for preparing compounds of the formula I as claimed in claim

1, where  $R^5 = \text{SOR}^8$ ,  $\text{SO}_2\text{R}^8$ , which comprises reacting a compound of the formula I $\beta$  ( $\equiv$  I where  $R^5 = \text{SR}^8$ ),




I where  $R^5 = \text{SR}^8$

where the variables  $R^1$  to  $R^8$  and  $[1]_1$  are each as defined in claim 1, with an oxidizing agent.

Please add new claims 14-25 as follows

14.(newly added) A cyclohexenonequinolinoyl derivative of the formula I



I

where:

$R^1$  is hydrogen, nitro, halogen, cyano,  $\text{C}_1\text{-C}_6\text{-alkyl}$ ,  $\text{C}_1\text{-C}_6\text{-haloalkyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxyiminomethyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxy}$ ,  $\text{C}_1\text{-C}_6\text{-haloalkoxy}$ ,  $\text{C}_1\text{-C}_6\text{-alkylthio}$ ,  $\text{C}_1\text{-C}_6\text{-haloalkylthio}$ ,  $\text{C}_1\text{-C}_6\text{-alkylsulfinyl}$ ,  $\text{C}_1\text{-C}_6\text{-haloalkylsulfinyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkylsulfonyl}$ ,  $\text{C}_1\text{-C}_6\text{-haloalkylsulfonyl}$ , aminosulfonyl,  $\text{N-(C}_1\text{-C}_6\text{-alkyl)}$

alkyl)aminosulfonyl.

N, N-di-(C<sub>1</sub>-C<sub>6</sub>-alkyl) aminosulfonyl.

N-(C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl)amino.

N-(C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfonyl)amino.

N-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-N-(C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl)amino.

N-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-N-(C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfonyl)amino.


phenoxy, heterocycloxy, phenylthio or heterocyclithio, where the four last-mentioned radicals may be partially or fully halogenated and/or may carry one to three of the following substituents :

nitro, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl.

C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy;

R<sup>2</sup>, R<sup>3</sup> are hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl or halogen;

R<sup>4</sup> is a compound IIa



where

IIa

R<sup>5</sup> is halogen, OR<sup>7</sup>, SR<sup>7</sup>, SOR<sup>8</sup>, SO<sub>2</sub>R<sup>8</sup>, OSO<sub>2</sub>R<sup>8</sup>, POR<sup>8</sup>R<sup>9</sup>,

OPR<sup>8</sup>R<sup>9</sup>, OPOR<sup>8</sup>R<sup>9</sup>, OPSR<sup>8</sup>R<sup>9</sup>, NR<sup>10</sup>R<sup>11</sup>, ONR<sup>11</sup>R<sup>12</sup>, N-linked

heterocyclyl or O-(N-linked heterocyclyl), where the

heterocyclyl radical of the two last-mentioned substituents

may be partially or fully halogenated and/or may carry one to

three of the following radicals:



nitro, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy;

R<sup>7</sup> is C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-cyloalkyl, C<sub>1</sub>-C<sub>20</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkenylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkynylcarbonyl, C<sub>3</sub>-C<sub>6</sub>-cyloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>3</sub>-C<sub>6</sub>-alkenyloxy carbonyl, C<sub>3</sub>-C<sub>6</sub>-alkynyloxy carbonyl, (C<sub>1</sub>-C<sub>20</sub>-alkylthio)carbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>6</sub>-alkenylaminocarbonyl, C<sub>3</sub>-C<sub>6</sub>-alkynylaminocarbonyl, N,N-di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)aminocarbonyl, N-(C<sub>3</sub>-C<sub>6</sub>-alkenyl)-N-(C<sub>1</sub>-C<sub>6</sub>-alkyl) aminocarbonyl, N-(C<sub>3</sub>-C<sub>6</sub>-alkynyl)-N-(C<sub>1</sub>-C<sub>6</sub>-alkyl) aminocarbonyl, N-(C<sub>1</sub>-C<sub>6</sub>-alkoxy)-N-(C<sub>1</sub>-C<sub>6</sub>-alkyl) aminocarbonyl, N-(C<sub>3</sub>-C<sub>6</sub>-alkenyl)-N-(C<sub>1</sub>-C<sub>6</sub>-alkoxy) aminocarbonyl, N-(C<sub>3</sub>-C<sub>6</sub>-alkynyl)-N-(C<sub>1</sub>-C<sub>6</sub>-alkoxy) aminocarbonyl, di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-aminothiocarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxyimino-C<sub>1</sub>-C<sub>6</sub>-alkyl, N-(C<sub>1</sub>-C<sub>6</sub>-alkylamino) imino-C<sub>1</sub>-C<sub>6</sub>-alkyl or N,N-di-(C<sub>1</sub>-C<sub>6</sub>-alkylamino)imino-C<sub>1</sub>-C<sub>6</sub>-alkyl, where the above-mentioned alkyl, cycloalkyl and alkoxy radicals may be partially or fully halogenated and/or may carry one to three of the following groups: cyano, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino-C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, hydroxycarbonyl, C<sub>1</sub>-C<sub>4</sub>-

alkylaminocarbonyl, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl)aminocarbonyl, aminocarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyloxy or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl;  
phenyl, heterocyclyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, heterocyclyl-C<sub>1</sub>-C<sub>6</sub>-alkyl,  
phenylcarbonyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, heterocyclylcarbonyl-C<sub>1</sub>-C<sub>6</sub>-alkyl,  
phenylcarbonyl, heterocyclylcarbonyl, phenoxy, carbonyl,  
heterocycliloxy, carbonyl, phenoxythiocarbonyl,  
heterocycliloxythiocarbonyl, phenoxy-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl,  
heterocycliloxy-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, phenylaminocarbonyl, N-(C<sub>1</sub>-C<sub>6</sub>-  
alkyl)-N-(phenyl)aminocarbonyl, heterocyclylaminocarbonyl, N-(C<sub>1</sub>-C<sub>6</sub>-  
alkyl)-N-(heterocyclyl)aminocarbonyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkenylcarbonyl or  
heterocyclyl-C<sub>2</sub>-C<sub>6</sub>-alkenylcarbonyl, where the phenyl and the heterocyclyl  
radical of the 20 last-mentioned substituents may be partially or fully  
halogenated and/or may carry one to three of the following radicals:  
nitro, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-halogenalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-  
haloalkoxy;

R<sup>8</sup>, R<sup>9</sup> are C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-  
alkynyl, C<sub>3</sub>-C<sub>6</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, hydroxyl, C<sub>1</sub>-C<sub>6</sub>-  
alkoxy, amino, C<sub>1</sub>-C<sub>6</sub>-alkylamino, C<sub>1</sub>-C<sub>6</sub>-haloalkylamino, di-  
(C<sub>1</sub>-C<sub>6</sub>-alkyl) amino or di-(C<sub>1</sub>-C<sub>6</sub>-haloalkyl) amino, where the  
abovementioned alkyl, cycloalkyl and alkoxy radicals may be  
partially or fully halogenated and/or may carry one to three  
of the following groups:  
cyano, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl) amino,  
C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-  
C<sub>4</sub>-alkoxycarbonyl, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl) amino-C<sub>1</sub>-C<sub>4</sub>-  
alkoxycarbonyl,  
hydroxycarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, di-(C<sub>1</sub>-C<sub>4</sub>-  
alkyl)aminocarbonyl, aminocarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyloxy  
or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl;

phenyl, heterocyclyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, heterocyclyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, phenoxy, heterocycliloxy, where the phenyl and the heterocyclyl radical of the last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following radicals:

nitro, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy;

R<sup>10</sup> is C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, hydroxyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>3</sub>-C<sub>6</sub>-alkenyloxy, C<sub>3</sub>-C<sub>6</sub>-alkynyloxy, amino, C<sub>1</sub>-C<sub>6</sub>-alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)amino or C<sub>1</sub>-C<sub>6</sub>-alkylcarbonylamino, where the abovementioned alkyl, cycloalkyl and alkoxy radicals may be partially or fully halogenated and/or may carry one to three radicals from the following group:

cyano, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl)amino-C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, hydroxycarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, di-(C<sub>1</sub>-C<sub>4</sub>-alkyl)aminocarbonyl, aminocarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyloxy or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl;

phenyl, heterocyclyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl or heterocyclyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, where the phenyl or heterocyclyl radical of the four last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following radicals:

nitro, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy;

R<sup>11</sup>, R<sup>12</sup> are C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl or C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl;

I is 0;

and their agriculturally useful salts.

15.(newly added) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim 14, where

R<sup>1</sup> is halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkylthio, heterocycloxy or phenylthio, where the two last-mentioned radicals may be partially or fully halogenated and/or may carry one to three of the substituents mentioned below:

nitro, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy;

R<sup>5</sup> is halogen, OR<sup>7</sup>, SR<sup>7</sup>, SOR<sup>8</sup>, SO<sub>2</sub>R<sup>8</sup>, OSO<sub>2</sub>R<sup>8</sup>, OPR<sup>8</sup>R<sup>9</sup>, OPOR<sup>8</sup>R<sup>9</sup>, OPSR<sup>8</sup>R<sup>9</sup>, NR<sup>10</sup>R<sup>11</sup> or N-bonded heterocyclyl which may be partially or fully halogenated and/or may carry one to three of the following radicals: nitro, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy .

16.(newly added) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim 14, where

R<sup>5</sup> is halogen, OR<sup>7</sup>, NR<sup>10</sup>R<sup>11</sup> or N-bonded heterocyclyl which may be partially or fully halogenated and/or may carry one to three of the following radicals:

nitro, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy.

17.(newly added) A cyclohexenonequinolinoyl derivative of the formula I as claimed in claim 14, where

R<sup>7</sup> is C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>20</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, (C<sub>1</sub>-C<sub>20</sub>-alkylthio)carbonyl, N,N-di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)aminocarbonyl, phenyl, phenylcarbonyl or phenoxy-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, where the phenyl radical of the three last-mentioned substituents may be partially or fully halogenated and/or may carry one to

three of the following radicals:


nitro, cyano, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy or C<sub>1</sub>-

C<sub>4</sub>-haloalkoxy;

R<sup>10</sup> is C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>1</sub>-C<sub>6</sub>-alkoxy;


R<sup>11</sup> is C<sub>1</sub>-C<sub>6</sub>-alkyl.

- 18.(newly added) A process for preparing compounds of the formula I as claimed in claim 14 where R<sup>5</sup> = halogen, which comprises reacting a cyclohexanedione derivative of the formula III,




where the variables R<sup>1</sup> to R<sup>3</sup>, and I are each as defined in claim 14, with a halogenating agent.

- 19.(newly added) A process for preparing compounds of the formula I as claimed in claim 14 where R<sup>5</sup> = OR<sup>7</sup>, OSO<sub>2</sub>R<sup>8</sup>, OPR<sup>8</sup>R<sup>9</sup>, OPOR<sup>8</sup>R<sup>9</sup> or OPSR<sup>8</sup>R<sup>9</sup>, which comprises reacting a cyclohexanedione derivative of the formula III,




where the variables R<sup>1</sup> to R<sup>3</sup>, and I are each as defined in claim 14, with a compound of the formula IVα, IVβ, IVγ, IVδ or IVε,




where the variables R<sup>7</sup> to R<sup>9</sup> are each as defined in claim 14 and L<sup>1</sup> is a nucleophilically replaceable leaving group.

20. (Newly added) A process for preparing compounds of the formula I as claimed in claim 14 where  $R^5 = OR^7, SR^7, POR^8R^9, NR^{10}R^{11}, ONR^{11}R^{12}$ , N-linked heterocyclyl or O-(N-linked heterocyclyl), which comprises reacting a compound of the formula I  $\alpha$  ( $\equiv$  I where  $R^5 = \text{halogen, } OSO_2R^8$ ).




I where  $R^5 = \text{halogen or } OSO_2R^8$

where the variables  $R^1$  to  $R^3$ , and I are each as defined in claim 14, with a compound of the formula  $V\alpha, V\beta, Vy, V\delta, Ve, V\eta, V\theta$ .



where the variables  $R^7$  to  $R^{12}$  are each as defined in claim 14, if appropriate in the presence of a base.

21. (Newly added) A process for preparing compounds of the formula I as claimed in claim 14 where  $R^5 = SOR^8, SO_2R^8$ , which comprises reacting a compound of the formula I  $\beta$  ( $\equiv$  I where  $R^5 = SR^8$ ).



I where  $R^5 = SR^8$

where the variables R<sup>1</sup> to R<sup>5</sup>, R<sup>7</sup>, R<sup>8</sup> and I are each as defined in claim 14, with an oxidizing agent.

22.(newly added) A composition, comprising a herbicidally effective amount of at least one cyclohexenon-equinolinoyl derivative of the formula I or an agriculturally useful salt of formula I as claimed in claim 14 and auxiliaries which are customarily used for formulating crop protection agents.

23.(newly added) A process for preparing compositions as claimed in claim 22, which comprises mixing a herbicidally effective amount of at least one cyclohexenonequinolinoyl derivative of the formula I or an agriculturally useful salt of formula I and auxiliaries which are customarily used for formulating crop protection agents.

24.(newly added) A method for controlling undesirable vegetation, which comprises allowing a herbicidally effective amount of at least one cyclohexenonequinolinoyl derivative of the formula I or an agriculturally useful salt of formula I as claimed in claim 14 to act on plants, their habitat and/or on seeds.

25.(newly added) The use of cyclohexenonequinolinoyl derivatives of the formula I or their agriculturally useful salts as claimed in claim 14 as herbicides.